

Appendix 1

Shared and Complementary Indicators within the National Adult Social Care, NHS and Public Health Outcomes Frameworks

Indicator	ASC Outcomes Framework	NHS Outcomes Framework	CCG Outcomes Framework	PH Outcomes Framework	BCF Indicators
Social care related quality of life					
Health-related quality of life for people with long-term conditions					
Employment of people with long-term conditions					
People with long term conditions feeling supported to manage their condition					
Carer-reported quality of life					
Proportion of adults with a learning disability in paid employment					
Proportion of adults in contact with secondary mental health services in paid employment					
Proportion of adults with a learning disability who live in their own home or with their family					
Proportion of adults in contact with secondary mental health services living independently, with or without support					
Proportion of people who use services, and their carers, who reported that they had as much social contact as they would like					
Patient and service user experience of integrated care					

Indicator	ASC Outcomes Framework	NHS Outcomes Framework	CCG Outcomes Framework	PH Outcomes Framework	BCF Indicators
Delayed transfers of care from hospital, and those which are attributable to adult social care					
Emergency readmissions within 30 days of discharge from hospital (males)					
Emergency readmissions within 30 days of discharge from hospital (females)					
Avoidable emergency admissions					
Proportion of older people offered rehabilitation following discharge from an acute or community hospital					
Proportion of older people (65 and over) who were still at home 91 days after discharge from hospital into reablement / rehabilitation services					
Admissions to long term care homes (people aged 18-64)					
Admissions to long term care homes (people aged 65+)					
Estimated diagnosis rate for people with dementia					
Dementia - The effectiveness of post-diagnosis care in sustaining independence and quality of life					
The proportion of people who use services who feel safe					
Under 75 mortality rate from cardiovascular disease					
Under 75 mortality rate from respiratory disease					
Under 75 mortality rate from liver disease					
Under 75 mortality rate from cancer					
Excess under 75 mortality rate in adults with serious mental illness					
Infant mortality					

